

On the non-closure of particle backscattering coefficient in oligotrophic oceans

ZhongPing Lee^{*1}, Yannick Huot²

¹ School For the Environment, University of Massachusetts Boston, 100 Morrissey Blvd. Boston, MA 02125-3393, USA

² Département de géomatique appliquée, Université de Sherbrooke, Sherbrooke QC, J1K 2R1 Canada
^{*}ZhongPing.Lee@umb.edu

Abstract: Many studies have consistently found that the particle backscattering coefficient (b_{bp}) in oligotrophic oceans estimated from remote-sensing reflectance (R_{rs}) using semi-analytical algorithms is higher than that from *in situ* measurements. This overestimation can be as high as ~300% for some oligotrophic ocean regions. Various sources potentially responsible for this discrepancy are examined. Further, after applying an empirical algorithm to correct the impact from Raman scattering, it is found that b_{bp} from analytical inversion of R_{rs} is in good agreement with that from *in situ* measurements, and that a closure is achieved.

©2014 Optical Society of America

OCIS codes: (010.4450) Oceanic optics; (280.0280) Remote sensing and sensors.

References and links

1. C. D. Mobley, *Light and Water: Radiative Transfer in Natural Waters* (Academic Press, 1994).
2. J. R. V. Zaneveld, "A theoretical derivation of the dependence of the remotely sensed reflectance of the ocean on the inherent optical properties," *J. Geophys. Res.* **100**(C7), 13135–13142 (1995).
3. R. W. Preisendorfer, *Hydrologic optics vol. 1: introduction* (National Technical Information Service. Also available on CD, Office of Naval Research, Springfield, 1976).
4. H. R. Gordon, "Spectral variations in the volume scattering function at large angles in natural waters," *J. Opt. Soc. Am.* **64**(6), 773–775 (1974).
5. M. Chami, D. McKee, E. Leymarie, and G. Khomenko, "Influence of the angular shape of the volume-scattering function and multiple scattering on remote sensing reflectance," *Appl. Opt.* **45**(36), 9210–9220 (2006).
6. H. R. Gordon, O. B. Brown, R. H. Evans, J. W. Brown, R. C. Smith, K. S. Baker, and D. K. Clark, "A semianalytic radiance model of ocean color," *J. Geophys. Res.* **93**(D9), 10909–10924 (1988).
7. A. Morel and B. Gentili, "Diffuse reflectance of oceanic waters. II Bidirectional aspects," *Appl. Opt.* **32**(33), 6864–6879 (1993).
8. S. Sugihara, M. Kishino, and N. Okami, "Contribution of Raman scattering to upward irradiance in the sea," *J. Oceanogr. Soc. Jpn* **40**(6), 397–404 (1984).
9. R. H. Stavn and A. D. Weidemann, "Optical modeling of clear ocean light fields: Raman scattering effects," *Appl. Opt.* **27**(19), 4002–4011 (1988).
10. H. R. Gordon, "Diffuse reflectance of the ocean: the theory of its augmentation by chlorophyll a fluorescence at 685 nm," *Appl. Opt.* **18**(8), 1161–1166 (1979).
11. S. K. Hawes, K. L. Carder, and G. R. Harvey, "Quantum fluorescence efficiencies of marine humic and fulvic acids: effects on ocean color and fluorometric detection," presented at the Ocean Optics, 1992.
12. Y.-H. Ahn, A. Bricaud, and A. Morel, "Light backscattering efficiency and related properties of some phytoplankters," *Deep-Sea Res.* **39**(11-12), 1835–1855 (1992).
13. D. Stramski, R. A. Reynolds, M. Kahru, and B. G. Mitchell, "Estimation of particulate organic carbon in the ocean from satellite remote sensing," *Science* **285**(5425), 239–242 (1999).
14. M. J. Behrenfeld, E. Boss, D. Siegel, and D. M. Shea, "Carbon-based ocean productivity and phytoplankton physiology from space," *Global Biogeochem. Cycles* **19**(1), GB1006 (2005), doi:10.1029/2004GB002299.
15. C. A. Brown, Y. Huot, P. J. Werdell, B. Gentili, and H. Claustre, "The origin and global distribution of second order variability in satellite ocean color and its potential applications to algorithm development," *Remote Sens. Environ.* **112**(12), 4186–4203 (2008).
16. IOCCG, "Remote Sensing of Inherent Optical Properties: Fundamentals, Tests of Algorithms, and Applications," in *Reports of the International Ocean-Colour Coordinating Group, No. 5, Z.-P. Lee, ed.* (IOCCG, Dartmouth, Canada, 2006), p. 126.
17. R. A. Maffione and D. R. Dana, "Instruments and methods for measuring the backward-scattering coefficient of ocean waters," *Appl. Opt.* **36**(24), 6057–6067 (1997).

18. IOCCG, "Remote Sensing of Ocean Colour in Coastal, and Other Optically-Complex, Waters," in *Reports of the International Ocean-Colour Coordinating Group, No.3*, S. Sathyendranath, ed. (IOCCG, Dartmouth, Canada, 2000).
19. A. H. Barnard, J. R. Zaneveld, and W. S. Pegau, "In situ determination of the remotely sensed reflectance and the absorption coefficient: closure and inversion," *Appl. Opt.* **38**(24), 5108–5117 (1999).
20. G. C. Chang, T. D. Dickey, C. D. Mobley, E. Boss, and W. S. Pegau, "Toward closure of upwelling radiance in coastal waters," *Appl. Opt.* **42**(9), 1574–1582 (2003).
21. D. M. O'Donnell, S. W. Effler, and G. A. Leshkevich, "Optical characterizations and pursuit of optical closure for the western basin of Lake Erie through in situ measurements," *Great Lakes Research* (2011).
22. H. R. Gordon, M. R. Lewis, S. D. McLean, M. S. Twardowski, S. A. Freeman, K. J. Voss, and G. C. Boynton, "Spectra of particulate backscattering in natural waters," *Opt. Express* **17**(18), 16192–16208 (2009).
23. Y. Huot, A. Morel, M. S. Twardowski, D. Stramski, and R. A. Reynolds, "Particle optical backscattering along a chlorophyll gradient in the upper layer of the eastern South Pacific Ocean," *Biogeosciences* **5**(2), 495–507 (2008).
24. R. J. W. Brewin, G. Dall'Olmo, S. Sathyendranath, and N. J. Hardman-Mountford, "Particle backscattering as a function of chlorophyll and phytoplankton size structure in the open-ocean," *Opt. Express* **20**(16), 17632–17652 (2012).
25. P. J. Werdell, B. A. Franz, S. W. Bailey, G. C. Feldman, E. Boss, V. E. Brando, M. Dowell, T. Hirata, S. J. Lavender, Z. Lee, H. Loisel, S. Maritorena, F. Mélin, T. S. Moore, T. J. Smyth, D. Antoine, E. Devred, O. H. F. d'Andon, and A. Mangin, "Generalized ocean color inversion model for retrieving marine inherent optical properties," *Appl. Opt.* **52**(10), 2019–2037 (2013).
26. J. Ras, H. Claustre, and J. Uitz, "Spatial Variability of Phytoplankton Pigment Distributions in the Subtropical South Pacific Ocean: Comparison Between in Situ and Predicted Data," *Biogeosciences* **5**(2), 353–369 (2008).
27. R. J. W. Brewin, S. Sathyendranath, D. Müller, C. Brockmann, P.-Y. Deschamps, E. Devred, R. Doerffer, N. Fomferra, B. Franz, M. Granta, S. Groom, A. Horseman, C. Hu, H. Krasemann, Z. Lee, S. Maritorena, F. Mélin, M. Peters, T. Platt, P. Regner, T. Smyth, F. Steinmetz, J. Swinton, J. Werdell, and G. N. White III, "The Ocean Colour Climate Change Initiative: A round-robin comparison on in-water bio-optical algorithms," *Deep-Sea Res.* (accepted).
28. Z.-P. Lee, R. Arnone, C. Hu, P. J. Werdell, and B. Lubac, "Uncertainties of optical parameters and their propagations in an analytical ocean color inversion algorithm," *Appl. Opt.* **49**(3), 369–381 (2010).
29. Z. P. Lee, K. L. Carder, and R. A. Arnone, "Deriving inherent optical properties from water color: a multiband quasi-analytical algorithm for optically deep waters," *Appl. Opt.* **41**(27), 5755–5772 (2002).
30. C. D. Mobley and L. K. Sundman, *HydroLight 5.2 User's Guide* (Sequoia Scientific, Inc., Bellevue, Washington., 2013).
31. G. Zibordi, F. Mélin, S. B. Hooker, D. D'Alimonte, and B. Holben, "An Autonomous Above-Water System for the Validation of Ocean Color Radiance Data," *IEEE Trans. Geosci. Rem. Sens.* **43**, 2 (2004).
32. G. Zibordi, J.-F. Berthon, F. Mélin, D. D'Alimonte, and S. Kaitala, "Validation of satellite ocean color primary products at optically complex coastal sites: Northern Adriatic Sea, Northern Baltic Proper and Gulf of Finland," *Remote Sens. Environ.* **113**(12), 2574–2591 (2009).
33. Z. P. Lee, K. L. Carder, C. D. Mobley, R. G. Steward, and J. S. Patch, "Hyperspectral remote sensing for shallow waters. I. A semianalytical model," *Appl. Opt.* **37**(27), 6329–6338 (1998).
34. Z.-P. Lee, K. Du, K. J. Voss, G. Zibordi, B. Lubac, R. Arnone, and A. Weidemann, "An inherent-optical-property-centered approach to correct the angular effects in water-leaving radiance," *Appl. Opt.* **50**(19), 3155–3167 (2011).
35. A. Morel, "Optical properties of pure water and pure sea water," in *Optical Aspects of Oceanography*, N. G. Jerlov, and Nielsen, E. S., ed. (Academic, New York, 1974), pp. 1–24.
36. X. Zhang, L. Hu, and M.-X. He, "Scattering by pure seawater: effect of salinity," *Opt. Express* **17**(7), 5698–5710 (2009).
37. A. Morel, B. Gentili, H. Claustre, A. Babin, A. Bricaud, J. Ras, and F. Tieche, "Optical properties of the "clearest" natural waters," *Limnol. Oceanogr.* **52**(1), 217–229 (2007).
38. N. B. Nelson and D. A. Siegel, "The global distribution and dynamics of chromophoric dissolved organic matter," *Annu. Rev. Mar. Sci.* **5**(1), 447–476 (2013), doi:10.1146/annurev-marine-120710-100751.
39. Y. Huot, C. A. Brown, and J. J. Cullen, "Retrieval of phytoplankton biomass from simultaneous inversion of reflectance, the diffuse attenuation coefficient, and Sun-induced fluorescence in coastal waters," *J. Geophys. Res.* **112**(C6), C06013 (2007), doi: 10.1029/2006JC003794.
40. A. Albert and C. D. Mobley, "An analytical model for subsurface irradiance and remote sensing reflectance in deep and shallow case-2 waters," *Opt. Express* **11**(22), 2873–2890 (2003).
41. Y.-J. Park and K. Ruddick, "Model of remote-sensing reflectance including bidirectional effects for case 1 and case 2 waters," *Appl. Opt.* **44**(7), 1236–1249 (2005).
42. Z. P. Lee, K. L. Carder, and K. P. Du, "Effects of molecular and particle scatterings on the model parameter for remote-sensing reflectance," *Appl. Opt.* **43**(25), 4957–4964 (2004).
43. J. M. Sullivan and M. S. Twardowski, "Angular shape of the oceanic particulate volume scattering function in the backward direction," *Appl. Opt.* **48**(35), 6811–6819 (2009).

44. D. Antoine, D. A. Siegel, T. Kostadinov, S. Maritorena, N. B. Nelson, B. Gentili, V. Vellucci, and N. Guillocheau, "Variability in Optical Particle Backscattering in Contrasting Bio-optical Oceanic Regimes," *Limnol. Oceanogr.* **56**(3), 955–973 (2011), doi:10.4319/lo.2011.4356.4313.0955.
45. Z. Lee, K. L. Carder, S. K. Hawes, R. G. Steward, T. G. Peacock, and C. O. Davis, "Model for the interpretation of hyperspectral remote-sensing reflectance," *Appl. Opt.* **33**(24), 5721–5732 (1994).
46. H. R. Gordon, "Contribution of Raman scattering to water-leaving radiance: a reexamination," *Appl. Opt.* **38**(15), 3166–3174 (1999).
47. T. K. Westberry, E. Boss, and Z. Lee, "The Influence of Raman Scattering on Ocean Color Inversion Models," *Appl. Opt.*, under revision (2013).
48. H. Loisel and D. Stramski, "Estimation of the inherent optical properties of natural waters from the irradiance attenuation coefficient and reflectance in the presence of Raman scattering," *Appl. Opt.* **39**(18), 3001–3011 (2000).
49. P. J. Werdell and S. W. Bailey, "An improved bio-optical data set for ocean color algorithm development and satellite data product validation," *Remote Sens. Environ.* **98**(1), 122–140 (2005).
50. Z. Lee, C. Hu, S. Shang, K. Du, M. Lewis, R. Arnone, and R. Brewin, "Penetration of UV-Visible solar light in the global oceans: Insights from ocean color remote sensing," *J. Geophys. Res.* **118**(9), 4241–4255 (2013), doi:10.1002/jgrc.20308.
51. M. S. Twardowski, H. Claustre, S. A. Freeman, D. Stramski, and Y. Huot, "Optical backscattering properties of the "clearest" natural waters," *Biogeosciences* **4**(6), 1041–1058 (2007).
52. R. M. Pope and E. S. Fry, "Absorption spectrum (380–700 nm) of pure water. II. Integrating cavity measurements," *Appl. Opt.* **36**(33), 8710–8723 (1997).
53. R. C. Smith and K. S. Baker, "Optical properties of the clearest natural waters (200–800 nm)," *Appl. Opt.* **20**(2), 177–184 (1981).
54. D. Stramski, R. A. Reynolds, M. Babin, S. Kaczmarek, M. R. Lewis, R. Röttgers, A. Sciandra, M. Stramska, M. S. Twardowski, B. A. Franz, and H. Claustre, "Relationships between the surface concentration of particulate organic carbon and optical properties in the eastern South Pacific and eastern Atlantic Oceans," *Biogeosciences* **5**(1), 171–201 (2008).
55. Z.-P. Lee, S. Shang, C. Hu, M. Lewis, R. Arnone, Y. Li, and B. Lubac, "Time series of bio-optical properties in a subtropical gyre: Implications for the evaluation of inter-annual trends of biogeochemical properties," *J. Geophys. Res.* **115**(C9), C09012 (2010).
56. J. Sullivan, M. Twardowski, J. R. V. Zaneveld, and C. Moore, "Measuring optical backscattering in water," in *Light Scattering Reviews 7: Radiative Transfer and Optical Properties of Atmosphere and Underlying Surface*, A. Kokhanovsky, ed. (Springer Praxis Books, 2013), pp. 189–224.
57. E. Boss and W. S. Pegau, "Relationship of light scattering at an angle in the backward direction to the backscattering coefficient," *Appl. Opt.* **40**(30), 5503–5507 (2001).
58. M. Chami, E. Marken, J. J. Starnes, G. Khomenko, and G. Korotaev, "Variability of the relationship between the particulate backscattering coefficient and the volume scattering function measured at fixed angles," *J. Geophys. Res.* **111**(C5), C05013 (2006), doi:10.1029/2005JC003230.
59. T. Oishi, Y. Takahashi, A. Tanaka, M. Kishino, and A. Tsuchiya, "Relation between the backward- as well as total scattering coefficients and the volume scattering functions by cultured phytoplankton," *J. School Mar. Sci. Technol. Tokai Univ.* **53**, 1–15 (2002).
60. N. T. Briggs, W. H. Slade, E. Boss, and M. J. Perry, "Method for estimating mean particle size from high-frequency fluctuations in beam attenuation or scattering measurements," *Appl. Opt.* **52**(27), 6710–6725 (2013).
61. D. Stramski, A. Bricaud, and A. Morel, "Modeling the inherent optical properties of the ocean based on the detailed composition of the planktonic community," *Appl. Opt.* **40**(18), 2929–2945 (2001).

1. Introduction

The spectrum and amplitude of the light leaving the ocean depends on the constituents therein and therefore can be used to obtain information about these constituents. For remote sensing application, the water-leaving radiance (L_w , $\text{W m}^{-2} \text{sr}^{-1}$) is used to describe this light, and because L_w is proportional to the incoming light on the ocean, described by the downwelling irradiance (E_d , W m^{-2}), E_d is used to normalize L_w and remove the dependence of variable incident irradiance. The ratio L_w/E_d , termed the remote sensing reflectance (R_{rs} , sr^{-1}) [1], depends on the concentration of optically active constituents in the water and the geometry of observation (angular distribution of the light field and observation direction of the sensor).

The impact of the optically active constituents on R_{rs} can be described by inherent optical properties (IOPs) [2]. The IOPs depend only on the type and concentration of constituents and not on the ambient light field [3]. These IOPs are the absorption coefficient (a , m^{-1}) which describes the loss of photon by absorption, and the volume scattering function (VSF, $\text{m}^{-1} \text{sr}^{-1}$) which represents the angular distribution of scattered photons. Because photons leaving the

ocean are incoming from the atmosphere and returning to the atmosphere (i.e., returning backward), the total backscattering coefficient (b_b , m^{-1}), which is the integration of the VSF in the backward direction, is commonly used to parameterize the impact of the VSF on R_{rs} [4,5]. Such that a function of the form $R_{rs} = f(a, b_b)$ is generally used to describe the relationship between a , b_b and R_{rs} in the ocean [6,7]. This function will also have to describe the observation geometry, which can be approximated at first order by the satellite viewing angle and solar angle. Most of these variables have a spectral dependence. In addition to a and b_b , other processes contribute to the R_{rs} , and these processes are trans-spectral and will transfer energy from shorter to longer wavelengths, which lead to a larger amount of light in the longer wavelengths than would be expected solely from the absorption and backscattering processes [6,7]. One such trans-spectral process is Raman scattering [8,9], in which a fraction of the photons incident on water at a shorter wavelength lead to emission at a longer wavelength (with a constant wavenumber difference from the incident photon). Another trans-spectral process is fluorescence where photons are absorbed by a molecule and reemitted at longer wavelengths. The main constituents in ocean waters that fluoresce are chlorophyll (emission light is centered near 685 nm) [10] and CDOM with a broad emission band throughout the visible [11].

The particulate backscattering coefficient (b_{bp} , m^{-1}), the fraction of the total backscattering coefficient that is not due to water molecules, is an important IOP. Not only does it describe how light is returned to the ocean surface and thus forms the signal collected by satellite sensors, but its value, when interpreted in terms of particle population, has also been proposed to infer concentrations of organic and inorganic constituents in the oceans [12–15]. Because of this importance, both remote sensing [13,16] and *in situ* methods [17] have been developed to obtain measurements of b_{bp} in surface waters in the global oceans.

Remote sensing methods to retrieve b_{bp} are based on its relationship to R_{rs} . For empirical algorithms [13], functional relationships are described based on concurrent measurements of b_{bp} and R_{rs} . For such empirical algorithms, there is no involvement of the radiative transfer equation (RTE), where all the information required for the algorithms is provided by the measurements and consequently any measurement bias are directly reflected in the output. Empirical algorithms will also include all relevant trans-spectral effects as they covary with the input R_{rs} and do not require knowledge of the optical properties of waters. Semi-analytical methods [16,18], on the other hand, obtain b_{bp} based on approximations of relationships found through the radiative transfer theory between R_{rs} and the IOPs and employ a few optical and bio-optical properties and/or relationships. Thus, b_{bp} derived semi-analytically from R_{rs} is independent of measured b_{bp} *in situ* but depends on using an accurate approximation of the relationship between R_{rs} , IOPs and trans-spectral processes.

If every step necessary for both determinations is well controlled and understood, b_{bp} from both methods should agree with each other, i.e. there should be a closure [19–21], as demonstrated in Gordon et al. [22]. But the study [22] covered just two cases, and the inversion scheme uses vertical profiles of upwelling radiance and downwelling irradiance as inputs, which is very different from remote sensing inversion. Empirical approaches also indicated good matches [13,15], but such results do not indicate a closure between two independent determinations. Various studies, in particular those using R_{rs} as the input to analytically retrieve b_{bp} [23–25], however, have found that b_{bp} values derived from semi-analytical algorithms are generally higher than those estimated from *in situ* measurements, i.e. there is no closure on b_{bp} . For instance, it was found that b_{bp} derived from MODIS-Aqua R_{rs} is systematically higher than b_{bp} inferred from chlorophyll-a concentration ($[\text{Chl}]$, mg/m^3) by a factor of 2-3 (see Fig. A1 in Huot et al. [23] and Fig. 11 in Brown et al. [15]), where the relationship between b_{bp} and $[\text{Chl}]$ was developed based on concurrent measurements of $[\text{Chl}]$ obtained by HPLC and b_{bp} obtained by a BB3 sensor (Wetlabs, Inc.) [26]. Beyond the study by Huot et al. [23], the R_{rs} -derived b_{bp} in oligotrophic waters via spectral optimization

algorithms is also found much higher (up to a factor of 2) than that measured *in situ* in recent evaluations of algorithm performances [25,27].

In principle, the uncertainty in the b_{bp} derived analytically from R_{rs} are small at least for oligotrophic waters [28] where the absorption coefficients in the longer wavelengths (550 nm and longer) are nearly constant (see more detailed discussions below). This is well demonstrated when the semi-analytical algorithms [25,29] were applied to a data set [16] obtained from radiative transfer simulations with the Hydrolight software³⁰ when trans-spectral effects were not included. Comparison showed that the retrieved b_{bp} generally contains errors ~20% or less for oligotrophic conditions [16,25]. Errors in measured R_{rs} will certainly contribute to analytically derived b_{bp} , but the errors in measured R_{rs} are not systematic [31,32], therefore we should not expect systematically and significantly higher b_{bp} from errors in R_{rs} . It is necessary to address such large systematic overestimations of b_{bp} . And, more importantly, it is necessary to demonstrate if a closure can be achieved after these sources of error are taken into consideration. Here, we focus on a semi-analytical algorithm, the quasi-analytical algorithm (QAA) developed by Lee et al. [29], to examine these discrepancies. After rapidly reviewing the algorithm, we further address the sources that could contribute to the non-closure in b_{bp} , and then followed with evaluations of ocean color b_{bp} products after the sources of uncertainties/errors are considered.

2. Brief description of the QAA

For the retrieval of inherent optical properties (IOPs) from spectral R_{rs} , a quasi-analytical algorithm (QAA) was developed by Lee et al. [29]. QAA uses the relationship between IOPs and R_{rs} derived from the RTE, and makes very few assumptions in the process of algebraically solving for b_{bp} from R_{rs} . The uncertainty in QAA-derived b_{bp} , at least for oligotrophic waters, is small and well understood.

From the RTE, a simplified relationship between IOPs and reflectance derived from radiative transfer computations, where the trans-spectral effects are omitted, is [2,6]

$$r_{rs}(\lambda) = \left(g_0 + g_1 \frac{b_b(\lambda)}{a(\lambda) + b_b(\lambda)} \right) \frac{b_b(\lambda)}{a(\lambda) + b_b(\lambda)}. \quad (1)$$

Here r_{rs} is the subsurface remote-sensing reflectance, which has an uncertainty ~10% [6,33]. The coefficients g_0 and g_1 are wavelength-independent model parameters, which depend on sun-sensor angular geometry as well as the shape of particle phase functions [7,34]. b_b can be expressed as

$$b_b(\lambda) = b_{bw}(\lambda) + b_{bp}(\lambda), \quad (2)$$

with b_{bw} representing the backscattering contribution from water molecules, and is taken as a constant for the global oceans [35,36], although b_{bw} varies slightly with temperature and salinity [36]. Therefore, when b_b is known, it is straightforward to derive b_{bp} using Eq. (2).

As Eq. (1) indicates, once g_0 and g_1 are known, that there are just two unknowns for any given r_{rs} : a and b_b , so b_b can be easily derived when a is known. In the QAA scheme, a at a reference wavelength (λ_0) is estimated with

$$a(\lambda_0) = a_w(\lambda_0) + \Delta a(\lambda_0), \quad (3)$$

with $a_w(\lambda_0)$ the contribution of water molecules, and $\Delta a(\lambda_0)$ estimated empirically from the R_{rs} spectrum. For the oceanic waters, λ_0 is set as 550 nm for MODIS and 555 nm for SeaWiFS, where R_{rs} at these wavelengths are available, respectively.

The first step of the algorithm consists in estimating $a(\lambda_0)$ using an empirical algorithm based on R_{rs} band ratios to estimate $\Delta a(\lambda_0)$. As described in Lee et al. [29], for oligotrophic waters $a(\lambda_0)$ is dominated (>95%) by $a_w(\lambda_0)$, thus $a(\lambda_0)$ can be accurately estimated; errors in

estimating $\Delta a(\lambda_0)$ have negligible impact on $a(\lambda_0)$ for such waters. This is particularly true for the oceanic gyres, where the amount of optically active constituents other than water (such as phytoplankton and gelbstoff) are scarce [37,38]. Therefore, as long as $R_{rs}(550)$ (or $R_{rs}(555)$) is measured accurately and Eq. (1) is valid, $b_b(550)$ (or $b_b(555)$) can be estimated accurately. If b_{bw} is accurately known, or at the very least is the same value as that used when processing the *in situ* validation measurements, $b_{bp}(550)$ (or $b_{bp}(555)$) estimated from R_{rs} should match that from *in situ* measurements.

3. Impact of errors in the QAA components on the b_{bp} retrieval

As mentioned above, the QAA algorithm when applied to data simulated by the radiative transfer equation retrieves b_{bp} with very low errors [16], as is expected in waters where the optical properties are consistent with those of oligotrophic waters. Because QAA is essentially based on such simulations, this highlights that the algorithm structure and approximations allow accurate retrieval of b_{bp} for such conditions. The discrepancy found when compared with *in situ* measurements must thus originate from one of three sources. The first is that the radiative transfer simulations are not representing accurately the relationships between R_{rs} and IOPs *in situ* as expressed in Eq. (1) (and those that allow going from r_{rs} to R_{rs}). The second is that the optical properties of pure saltwater, used in Eqs. (2) and (3), are incorrect. The third is that there are errors or biases present in measured b_{bp} . We now examine these different potential sources of variability.

3.1 R_{rs} -IOP relationship

The relationship between R_{rs} and $b_b/(a + b_b)$ as described by Eq. (1) is not exact and other mathematical models were also developed in the past decades [39–42]. These expressions are approximate solutions of the RTE and generally ignored the trans-spectral effects. In addition, the use of b_b conveniently removes the complex relationship between r_{rs} and the volume scattering function, although a remote sensor measures radiance in one direction rather than the entire backscattered domain that is required to characterize b_b . For a hypothetical ocean without trans-spectral effects, this simplification results in errors up to ~12% compared with the simulations [6]. Depending on inputs used in data simulations (e.g., phase function of particle scattering, range of IOPs, etc.), slightly different g_0 and g_1 values are obtained. By changing the values of g_0 and g_1 using two sets of published values (i.e. from 0.0949 and 0.0794 [6] to 0.089 and 0.125 [29]), it is found that the retrieved b_{bp} values with the latter coefficients are higher than that with the former, and this difference is higher for smaller b_{bp} (see Fig. 1), with the maximum difference around 12% (for $b_{bp}(555) \sim 0.0005 \text{ m}^{-1}$). This analysis shows that the selection of values to represent g_0 and g_1 cannot be the reason leading to ~200-300% difference between R_{rs} -retrieved b_{bp} and *in situ* measured b_{bp} . This is further supported by the fact that in more eutrophic waters the retrieved and measured b_{bp} are much closer; therefore, for errors in g_0 and g_1 to be the source of the significant differences in b_{bp} observed it would require substantial changes in g_0 and g_1 with trophic status. The values of g_0 and g_1 vary mostly due to the geometry of observation (including angular distribution of the incident light field) and angular shape of the particle phase function in the backward domain [5,34]. While the former does not change with trophic status, observations [23,43,44] suggest that the latter also does not change strongly across trophic states.

A major caveat of Eq. (1) in modeling R_{rs} is the omission of contributions from in-elastic processes, which include those from chlorophyll *a* and CDOM fluorescence and Raman scattering [1]. Chlorophyll *a* fluorescence is centered around 685 nm (with a roughly Gaussian shape at a half-height width around 25 nm), so its influence in the QAA-retrieved b_{bp} is nil as QAA uses measurements around 550 nm to retrieve b_{bp} . For oceanic waters, the contribution from the CDOM fluorescence is small, so the impact of this effect is also negligible [11,45]. However, the contribution from the Raman scattering to R_{rs} can be as high as 20% in the longer wavelengths for oceanic waters [8,9,46], thus the impact of Raman

scattering on the retrieval of b_{bp} could be significant for clear oceanic waters [47,46,48]. Indeed for simulated waters with Raman contribution so extreme that $b_{bp} = 0$ and an absorption consistent with a chlorophyll concentration of 0.01 mg/m^3 , the QAA scheme returns a $b_{bp}(550)$ value of $\sim 0.00025 \text{ m}^{-1}$ as it interprets the light from Raman scattering to particulate backscattering 22; this value is approximately half of the lowest values of $b_{bp}(550)$ measured in the NOMAD data set [49].




Fig. 1. Impact of the g_0 and g_1 coefficients on the retrieval of $b_{bp}(555)$ via QAA, with data from the NOMAD data set. The $b_{bp}(555, \text{QAA})$ used g_0 and g_1 values as 0.089 and 0.125 respectively which correspond to the values in QAA [29] for sun at zenith and observation at nadir, while $b_{bp}(555, \text{Gordon})$ used g_0 and g_1 values as 0.0949 and 0.0794, respectively, as reported in Gordon et al. [6].

To effectively process ocean color satellite imagery, an empirical algorithm has been developed to correct this contribution in R_{rs} inversion [50], and Fig. 2 presents b_{bp} retrievals before and after the correction of Raman contribution for measurements made in the South Pacific gyre. For these “clearest” water of the global oceans [37,51], the ratio of R_{rs} -derived b_{bp} to *in situ* b_{bp} decreased from $\sim 300\%$ to $\sim 200\%$ (for lowest $b_{bp}(555)$), removing on average 0.00022 m^{-1} of retrieved $b_{bp}(555)$ from R_{rs} , clearly echoing the importance of removing Raman effect in the analytical retrieval of b_{bp} from R_{rs} [22,46–48]. As presented, however, b_{bp} from R_{rs} is still generally higher than that from *in situ* measurement, and this difference is $\sim 0.00025 \text{ m}^{-1}$ for all samples.

The scheme to empirically correct the Raman effect has a maximum error of about 15% in estimating the Raman contribution to R_{rs} [50]. The impact of a 15% adjustment in the Raman correction, however, results in just $\sim 3\%$ uncertainty in the R_{rs} derived b_{bp} . This indicates that the empirical scheme is sufficient for the correction of Raman scattering contribution in oceanic waters and that the origin of the remaining differences cannot originate from incorrect Raman correction.




Fig. 2. Ratio of Rrs-derived $b_{bp}(555)$ to *in situ* measured $b_{bp}(550)$ for waters in the South Pacific Gyre, before (open green) and after (solid blue) correcting the Raman contributions in the R_{rs} inversion process. *In situ* $b_{bp}(550)$ was determined by the Hydroscat sensor (HOBILabs) as described in Stramski et al.⁵⁴ while the *in situ* R_{rs} measurements were obtained using a modified HyperPro profiling radiometer (Satlantic Inc.) and a reference surface sensor and are described in Lee et al. [55]. There could be slight differences between $b_{bp}(550)$ and $b_{bp}(555)$ and this difference is ignored here.

3.2 Pure-water absorption coefficient

Another source of error in the QAA-derived b_{bp} is the uncertainty in the absorption coefficient of pure water at the reference wavelength (550 or 555 nm), since this value is used in the estimation of $a(\lambda_0)$ before the analytical derivation of $b_{bp}(\lambda_0)$. Based on Eq. (1), higher $a(555)$ will result in higher $b_b(555)$ and consequently higher $b_{bp}(555)$. Modern laboratory and field measurements (see a list of a_w values in Mobley¹) of $a_w(555)$ span approximately $0.0596 m^{-1}$ [52] to $0.0673 m^{-1}$ (spectrally interpolated from Smith and Baker [53]). QAA uses the lowest published a_w values [52], leading to the lowest values of $b_{bp}(555)$. Pope and Fry [52] also provide the standard deviation of the error on their measurement at 555 nm as $0.0012 m^{-1}$, this translates in QAA in an error of $\sim 4\%$ on the determination of $b_{bp}(555)$ for the data shown in Fig. 2. Obtaining optically pure water is very challenging and the value published may still suffer from minor contamination by absorbing components (as was the case with previous determinations). However, above ~ 450 nm [37], the $a_w(555)$ values of Pope and Fry [52] are consistent with values obtained from the diffuse attenuation coefficient of the clearest water which imposes a strong upper limit on the a_w value and strongly support the assertion that these are nearly devoid contribution from colored substance at 555 nm.

4. Uncertainties in b_{bp} measured *in situ*

The non-closure of b_{bp} does not have to originate only from errors or uncertainties in R_{rs} inversion; it also includes errors from *in situ* measurements [56]. While significant efforts were deployed to obtain accurate values in the clearest Pacific waters, these measurements are extremely challenging and push the sensitivity and resolution of the instruments to their limit.

Since the advent of commercial instruments, measurements of b_{bp} *in situ* are generally made with active instruments (e.g., Hydroscat, HOBILab, Inc.; BB3, WET Labs, Inc.). In these sensors, light from a pulsed LED is emitted into the water and the backscattered light (usually with a scattering angle of $\sim 120^\circ$ to 140°) from this source is recorded, and this

energy is converted to b_{bp} after subtracting the contribution of pure saltwater. Calibration of this system is usually carried out in the lab with well-characterized materials (reflecting surface or beads) [51]. Apart from possible calibration errors, there are two main sources of errors that can arise from these measurements. The first is caused by an estimation of b_{bp} from a measurement of the volume scattering function at 120° or 140°, while the other may arise from the undersampling of the backscattering from larger and less abundant particles due to the limited observation volumes.

The error caused by the estimation of b_{bp} from a measurement of the VSF between 120° to 140° (a constant factor χ is used for this conversion) has been studied by different authors [57–59]. All authors report that errors caused by this approximation should be less than ~10%. This conclusion arises in all cases from measured VSF in coastal waters as well as simulated VSF from Mie scattering. Although a thorough study of the conversion factor hasn't been carried out in oligotrophic ocean waters it appears unlikely that the error could increase by a factor of 10 and could explain the discrepancies observed. Therefore, the measurement strategy at a single angle does not seem to be a plausible source of error to explain the differences observed.

The passive measurement systems have a sampling volume in the range of 10s to 1,000s m^3 , while the instruments measuring b_{bp} *in situ* have a sampling volume around $10^{-6} m^3$, i.e. there is a many orders of magnitude difference in the sampling volume between the active backscatter sensors and the passive remote systems. In oceanic waters, larger particles are generally scarcer, thus the sampling volume of an active sensor will likely miss some of the larger particles during individual observations. This is demonstrated in Briggs et al. [60] where the observed spikes in b_b time series are indicators of large particles, although larger particles generally have a smaller backscattering efficiency [61]. The difference in sampling volume may help to explain the slightly ($\sim 0.0003 m^{-1}$) higher b_{bp} by Hydrosat than that by BB3 in the clearest waters, although both sensors were carefully calibrated for the field measurements (see Stramski et al. [54,51] for details). This is because Hydrosat has larger sampling volume than BB3 (or BB9) due to measurement geometry, thus more particles might be “counted” for by a Hydrosat sensor, although Hydrosat will not be able to cover all particles in the bulk water during a measurement scan. Indeed, if Hydrosat underestimated $b_{bp}(555)$ by $0.0002 m^{-1}$ (a very small number) [54], much better closure of b_{bp} is achieved between R_{rs} inversion and *in situ* measurements for waters in the South Pacific gyre (see Fig. 3), with the ratio of R_{rs} -derived $b_{bp}(555)$ to the adjusted (adding $0.0002 m^{-1}$ to the Hydrosat measured value) *in situ* $b_{bp}(555)$ in a range of ~ 0.8 - 1.2 (which was ~ 1.1 - 2.2 , see Fig. 2). The $\pm 20\%$ difference (transfers to $\sim 0.0001 m^{-1}$ absolute difference) is beyond the precision of either remote sensing measurements or *in situ* determinations. Indeed, even the value of $0.0002 m^{-1}$ that we added is within the instrument offset uncertainty (specified as between 5×10^{-5} to $5 \times 10^{-4} m^{-1}$, <http://www.hobilabs.com/>). These results suggest that it is necessary to take appropriate average over a long time or wide spatial range of *in situ* b_b measurements to have a “ground truth” to match the large volume observed by remote sensing systems. In particular, if the objective is to compare products of backscattering sensors with that from passive sensors, whether remotely or *in situ*, the removal of “spikes” must be carried out only if the spikes are caused by instrument malfunction and not by rare large particles (see user manual of the backscattering sensors).




Fig. 3. Ratio of Rrs-derived $b_{bp}(555)$ to *in situ* measured $b_{bp}(555)$ for waters in the South Pacific Gyre, with Rrs-derived $b_{bp}(555)$ after correcting the Raman contributions in the R_{rs} inversion process, and Hydroscat-measured $b_{bp}(555)$ was increased by 0.0002 m^{-1} from the original measurements.

To further examine the closure of b_{bp} on a global scale, we carried out an analysis similar as Huot et al. [23], where we compared the MODIS derived b_{bp} using QAA with *in situ* statistical relationships that have been derived (Fig. 4). As opposed to Huot et al. [23] the comparison is done at 547 nm instead of 443 nm to avoid spectral errors when extrapolating the value from 547 nm to 443 nm; furthermore, care was taken to use the same value of the backscattering coefficient of pure saltwater (though at constant salinity) for the remote sensing estimates as was used for the processing of the Hydroscat data underlying the Huot et al. [23] relationships. Before the Raman corrections there is clearly an overestimation of $b_{bp}(547)$ (Fig. 4(a), blue line vs pink line); this overestimation disappears after the correction, and the running median of the remote sensing data follows very closely the global statistical relationship derived by Brewin et al. [24]. These results confirm that the Raman effect is a big contributor to the non-closure of b_{bp} in previous studies and that the empirical scheme to correct this effect is plausible [23].

5. Conclusions

The results of this study highlighted the two major sources that contributed to the non-closure of b_{bp} in oligotrophic oceans shown in previous studies. One is the Raman scattering, which resulted in an increased b_{bp} when it is analytically derived from R_{rs} when the Raman effect was not considered; another is associated with the uncertainty in *in situ* measured b_{bp} (see Table 1 for summary). In particular, it is extremely difficult to get the “ground truth” of b_{bp} of such clear waters, as the values of b_{bp} are generally smaller than 0.001 m^{-1} . A closure in b_{bp} for oligotrophic water is achieved after correcting the Raman contribution in the inversion process and using the adjusted b_{bp} values (adding 0.0002 m^{-1} to that measured by Hydroscat, a value well within the measurement uncertainty). This closure in b_{bp} demonstrated the effectiveness of the empirical scheme in correcting the Raman scattering contribution, and provided a confidence in satellite ocean-color b_{bp} products, as long as all relevant processes are considered in its analytical inversion.




Fig. 4. Comparison of $b_{bp}(547)$ from QAA before (panel A) and after (panel B) Raman correction. The colored points represent the number of $b_{bp}(547)$ observations obtained from the QAA algorithm (see color bar) when applied on a global Level 3 (SMI) data set from the MODIS sensor for March 2007 (all daily level 3 values are temporally averaged). The Magenta lines (see legend in Panel A) are for three fits (i.e., with different values of the “Beta parameter”, see Huot et al. [23]) to three b_{bp} data sets obtained during the BIOSOPE cruise: “adopted” stands for the Hydroscat and BB3 data set, “Hydroscat Beta” stands for the Hydroscat data only and “BB3 Beta” stands for the BB3 data set only. The black continuous and dashed lines correspond to equation 6 and equation 14 respectively in Brewin et al. [24] as parameterized on their Dataset “A” at 526 nm; a λ^{-1} function was used to obtain b_{bp} at 547 nm. The blue line is a 1001 points running median on the MODIS data set.

Table 1. Summary of likely sources contributing to the non-closure of b_{bp}

Sources	Impact on estimate of b_{bp} with <i>in situ</i> sensor	Effect on estimate of $b_{bp}(550)$ from QAA
Chlorophyll fluorescence	Negligible except in the red wavebands near 685 nm.	nil
CDOM fluorescence	Negligible if any	Negligible for oligotrophic waters
Variability in the estimates of g_0 and g_1 parameters	N/A	~10% (this study), probably up to 15%. Variation of particle phase function could contribute a few more percent
Raman	N/A	Can contribute up to ~50% in estimated $b_{bp}(550)$. A bias of ~0.00025 m^{-1} is expected in very clear waters.
Inter instrument calibration and estimates	0.00025 – 0.0003 m^{-1} was observed in the South Pacific waters.	N/A
Instrumental offset uncertainty	0.00005 to 0.0005 m^{-1} (Hydroscat specs)	N/A
Estimate of b_{bp} from phase function at 120° or 140°	Less than ~10%.	N/A
Error on the absorption by pure water	Negligible if any	Smallest published value already

Acknowledgments

Financial support from the NASA Ocean Biology and Biogeochemistry and Water and Energy Cycle Programs and the JPSS VIIRS Ocean Color Cal/Val Project is greatly appreciated. We are grateful for the comments, and discussions, about b_b measurements from Drs. Jim Sullivan, Robert Brewin, and Giorgio Dall’Olmo. The acquisition of BIOSOPE data where funded through CNRS(1)-INSU(2) grants; we thank Marlon Lewis for sharing the Rrs data and Dariusz Stramski for sharing the Hydroscat data. Comments and suggestions from two anonymous reviewers significantly improved this article and are greatly appreciated.